

REMARK ON A CONJECTURE OF SHAFAREVICH

Robert Treger

We will assume several facts from [T, Sections 1, 2] and generalize one proposition from that note [T, Proposition 2.3] which is a special case of Shafarevich's conjecture (see [K, p. 9, (0.3.1)]).

Proposition. *Let $X \subset \mathbf{P}^r$ be a nonsingular connected complex n -dimensional projective variety with large residually finite fundamental group $\pi_1(X)$. We will also assume that a general curvilinear section through any point of X is not an elliptic curve. Then the universal covering of X , denoted by U , is a Stein manifold.*

Proof. Let $\mathcal{O}_X(m)$ be a fixed sufficiently very ample line bundle, i.e. $m \gg 0$, and $\mathcal{L} := \mathcal{O}_U(m)$ its pullback. As in [T, (2.2)], we can introduce a Hermitian metric on $\mathcal{O}_X(m)$ as well as a $\pi_1(X)$ -invariant Hermitian metric on $\mathcal{L}$. We obtain the corresponding section $B_{U,\mathcal{L}}(z, w)$ holomorphic in z and antiholomorphic in w . Thus $B_{U,\mathcal{L}}(z, w)$ is a Hermitian kernel which is not necessary positive, i.e., it may not satisfy the positivity condition [T, 1.4(ii)]. Let $\beta : U \rightarrow X$ be the covering map.

1.1. Given $U, \mathcal{L}$ and $B_{U,\mathcal{L}}(z, w)$ as above, we will show that U has a natural Kahler metric. Since $B(z, z) := B_{U,\mathcal{L}}(z, z) > 0$, we may consider $\log B(z, z)$. We will show that $\log B(z, z)$ is *strictly* plurisubharmonic, i.e.,

$$ds_U^2 = 2 \sum g_{jk} dz_j d\bar{z}_k \quad (g_{jk} := \frac{\partial^2}{\partial z_j \partial \bar{z}_k} \log B(z, z))$$

determines a Hermitian metric. For an arbitrary real tangent vector at $z \in U$,

$$\mathbf{v} = \partial_a + \bar{\partial}_a \quad (\partial_a := \sum_{t=1}^n a_t \frac{\partial}{\partial z_t}, \quad \bar{\partial}_a := \sum_{t=1}^n \bar{a}_t \frac{\partial}{\partial \bar{z}_t}),$$

we set $|\mathbf{v}|^2 := \sum g_{jk} a_j \bar{a}_k$ hence $|\mathbf{v}|^2 = \partial_a \bar{\partial}_a \log B(z, z)$. The g_{jk} 's determine a Hermitian metric iff $|\mathbf{v}|^2 > 0$ for $\mathbf{v} \neq 0$. With the above notation,

$$\begin{aligned} \partial_a \bar{\partial}_a \log B(z, z) &= \frac{1}{B(z, z)^2} \left\{ \sum_k |\mathbf{v} \psi_k(z)|^2 \cdot \sum_k |\psi_k(z)|^2 - \left| \sum_k \mathbf{v} \psi_k(z) \cdot \overline{\psi_k(z)} \right|^2 \right\} \\ &\geq \frac{1}{B(z, z)^2} \left\{ \sum_k |\mathbf{v} \psi_k(z)|^2 \cdot \sum_k |\psi_k(z)|^2 - \left(\sum_k |\mathbf{v} \psi_k(z) \cdot \overline{\psi_k(z)}| \right)^2 \right\} \end{aligned}$$

where $B(z, z) = \sum \psi_k(z) \overline{\psi_k(z)}$ in a neighborhood of z (the calculations are similar to [FK, pp. 13–14, pp. 190–191]). In the Lagrange identity

$$\sum_{k=1}^s A_k^2 \sum_{k=1}^s B_k^2 - \left(\sum_{k=1}^s A_k B_k \right)^2 = \frac{1}{2} \sum_{i=1}^s \sum_{j=1}^s (A_i B_j - B_i A_j)^2 \quad (\forall A_i, \forall B_i \in \mathbf{R}_+, s \in \mathbf{N}),$$

the right-hand side is strictly positive for $s = \infty$, provided it is strictly positive for some $s \in \mathbf{N}$ (and the series are convergent). It follows that $|\mathbf{v}|^2 > 0$ for $\mathbf{v} \neq 0$, and we get the desired metric.

1.2. Set $\Phi := \log B$. Let $\mathbf{p}$ and $\mathbf{q}$ be two points of U with coordinates $(z(\mathbf{p}))$ and $(z(\mathbf{q}))$. We assume $\mathbf{p}$ and $\mathbf{q}$ belong to a small open subset $\mathcal{V} \subset U$. The functional element of diastasis of the above metric is defined as follows (Calabi [C, (5)]):

$$D_U(\mathbf{p}, \mathbf{q}) := \Phi(z(\mathbf{p}), \overline{z(\mathbf{p})}) + \Phi(z(\mathbf{q}), \overline{z(\mathbf{q})}) - \Phi(z(\mathbf{p}), \overline{z(\mathbf{q})}) - \Phi(z(\mathbf{q}), \overline{z(\mathbf{p})}).$$

It is strictly plurisubharmonic in $z(\mathbf{q})$ for a fixed $\mathbf{p}$, and inductive on holomorphic submanifolds of a manifold with analytic Kahler metric [C, Propositions 4 and 6].

For a fixed $\mathbf{p} \in U$, $f(\mathbf{q}) := D_U(\mathbf{p}, \mathbf{q})$ is a *unique* primitive function, defined in some neighborhood of $\mathbf{p}$, that satisfies the vanishing of the corresponding partial derivatives [C, Chap. 2], [U, Appendix].

To conclude the proof of our proposition, it suffices to show that the $f(\mathbf{q})$ generates a single-valued strictly plurisubharmonic function $P_U(x)$ on U such that, for any real number c , the set $U_c := \{x \in U \mid P_U(x) < c\}$ is relatively compact in U . Then U will be Stein by the Oka-Grauert-Narasimhan solution of the Levi problem.

1.3. We will show that a functional element $f(\mathbf{q}) = D_U(\mathbf{p}, \mathbf{q})$ can be prolonged to a single-valued function along a path in U originating in $\mathbf{p}$ so that we get the corresponding functional element at each point of the path.

First, any holomorphic bundle on a noncompact Riemann surface is trivial. Second, given a finite number of points on U , as in [T, Remark 2.2.1] we can assume they lie on a connected noncompact Riemann surface $Y \subset U$ which is a Galois covering of a curvilinear section of X_γ ($\gamma \gg 0$, γ depends on the points) by Deligne's theorem [K, Theorem 2.14.1] since $\pi_1(X)$ is large. Thus, if a path $\alpha \subset Y$ then the functional element $f(\mathbf{q})$ can be prolonged along α .

Finally, for an arbitrary path $\alpha \subset U$ with initial point $\mathbf{p}_0$ and end point $\mathbf{p}_1$, one can choose an arbitrary finite number of points on α . As above, we obtain a connected noncompact Riemann surface $Y \subset U$ passing through $\mathbf{p}_0$, $\mathbf{p}_1$, and the chosen points. We get the desired prolongation along α . Since U is simply connected, the functional element $f(\mathbf{q})$ generates a function $P_U(x)$ on the whole U .

1.4. Since X is compact, to establish the relative compactness of U_c (see 1.2), it suffices to show $|U_c \cap \beta^{-1}(x)| < \infty$ for $\forall x \in X$. The inequality is trivial if $\dim X = 1$. Suppose to the contrary $|U_c \cap \beta^{-1}(x)| = \infty$ for some $x \in X$ with $\dim X > 1$.

We take a general curvilinear section C_x through x and obtain the covering $\nu : \Delta \rightarrow S$, where $S := \beta^{-1}(C_x)$ and Δ is a disk. The corresponding P_Δ (arising from the Poincaré-Bergman metric) goes to infinity on any infinite discrete subset in Δ . Furthermore, the relative Poincaré map transforms P_Δ into P_S . We derive a contradiction at once from the classical (integral) lifting formula for automorphic functions, i.e., P_S lifts to P_Δ and the relative Poincaré map transforms P_Δ back into P_S .

This concludes the proof of the proposition.

REFERENCES

- [C] E. Calabi, *Isometric imbedding of complex manifolds*, Ann. of Math. **58** (1953), 1–23.
- [FK] J. Faraut, S. Koneyuki, A. Korányi, Q.-k. Lu, G. Roos, *Analysis and geometry on complex homogeneous domains*, Birkhäuser, Boston, 2000.
- [K] J. Kollár, *Shafarevich maps and automorphic forms*, Princeton Univ. Press, Princeton, 1995.
- [T] R. Treger, *Uniformization*, arXiv:math.AG/1001.1951.
- [U] M. Umehara, *Kähler Submanifolds of Complex Space Forms*, Tokyo J. Math. **10** (1987), 203–214.

PRINCETON, NJ 08540

E-mail address: roberttreger117@gmail.com